IT'S TIME TO TALK MAVENCLAD

Find ways to talk with your healthcare provider to see if MAVENCLAD is right for you.

What is MAVENCLAD?

MAVENCLAD is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include relapsing-remitting disease and active secondary progressive disease, in adults. Because of its safety profile, MAVENCLAD is generally used in people who have tried another MS medicine that they could not tolerate or that has not worked well enough.

MAVENCLAD is not recommended for use in people with clinically isolated syndrome (CIS).

It is not known if MAVENCLAD is safe and effective in children under 18 years of age and is therefore not recommended.

MAVENCLAD may cause serious side effects.

Treatment with MAVENCLAD may increase your risk of developing cancer. You should follow healthcare provider instructions about screening for cancer. Because of the risk of birth defects, do not take MAVENCLAD if you are pregnant or of childbearing potential and not using effective birth control.

Please see additional Important Safety Information on pages 6-8 and MAVENCLAD Prescribing Information and Medication Guide, including **serious side effects**.

Questions? 1-877-447-3243

MAVENCLAD®
(cladribine) tablets 10 mg

IMPORTANT SAFETY INFORMATION

MAVENCLAD may cause serious side effects, including:

- **Risk of cancer (malignancies).** You should follow healthcare provider instructions about screening for cancer.
- **MAVENCLAD may cause birth defects if used during pregnancy.** Females must not be pregnant when they start treatment with MAVENCLAD or become pregnant during MAVENCLAD dosing and within 6 months after the last dose of each yearly treatment course. You should stop treatment with MAVENCLAD and contact your healthcare provider right away if you become pregnant during treatment with MAVENCLAD.

The person shown in this brochure is not an actual MAVENCLAD patient.

Please see additional Important Safety Information on pages 6-8 and MAVENCLAD Prescribing Information and Medication Guide, including **serious side effects**.

Questions? 1-877-447-3243

MAVENCLAD® (cladribine) tablets 10 mg

START THE CONVERSATION

To help you ask the right questions and get the right information from your healthcare provider, bring this guide to your next appointment and use it to get the conversation started.

You and your healthcare provider can decide if MAVENCLAD is right for you.

QUESTIONS ABOUT MAVENCLAD

It's important to have all the facts while considering a treatment option.
Be sure to ask your healthcare provider about:

Clinical Trial Results

Understanding how well MAVENCLAD works (efficacy) is important. Here are a few questions you may want to ask:

- Can MAVENCLAD be an effective treatment option for my relapsing MS?
- Has MAVENCLAD been shown to reduce the rate of relapses?
- Has MAVENCLAD been shown to slow disability progression?
- Has MAVENCLAD been shown to reduce the number of brain lesions?

Notes:

QUESTIONS ABOUT MAVENCLAD (CONTINUED)

Short-course Oral Dosing

Here are some questions to ask your healthcare provider about taking MAVENCLAD and its unique dosing schedule:

- What can you tell me about MAVENCLAD dosing?
- How is it different from other treatments?
- Once I start treatment, will I have to come back for any lab tests to monitor how I'm doing with MAVENCLAD?
- Since MAVENCLAD isn't taken every day, is it still working even when I'm not taking it?

Notes:

Safety Information

Here are some questions to ask your healthcare provider about MAVENCLAD and its safety:

- What side effects are associated with MAVENCLAD?
- Can you tell me about the risks I need to consider before treatment with MAVENCLAD?
- What were the cancer rates in clinical trials?
- Based on my medical history, are there any reasons why MAVENCLAD would not be the right option for me?

Notes:

OTHER QUESTIONS TO CONSIDER

- How does MAVENCLAD work?
- How long does MAVENCLAD stay in my system?
- When can I start considering pregnancy if I am treated with MAVENCLAD?

Notes:

GETTING STARTED WITH MAVENCLAD

Once you and your healthcare provider have chosen MAVENCLAD, it's time to make a treatment plan. Here are some questions to ask to get started:

- When can I begin treatment with MAVENCLAD?
- What's the process for starting?
- What kind of follow-up appointments and tests should I expect?
- Who can I speak to if I have questions or need support?
- (If currently on medication) When will I need to stop taking my current medication?

Notes:

MS LifeLines®

Think of MS LifeLines as a helping hand when it comes to living with relapsing MS. Sponsored by EMD Serono, MS LifeLines is a patient education support program that has been helping people with MS and their care partners for over 17 years.

MS LifeLines connects you with a team of MS-certified nurses and Financial Support Specialists. They've helped people in person and on over 4 million phone calls. MS LifeLines will be here for you throughout your treatment experience.

You can reach MS LifeLines at **1-877-447-3243**
Monday through Friday / 8 AM to 8 PM ET
Saturday / 9 AM to 5 PM ET

Questions? 1-877-447-3243

MAVENCLAD® (cladribine) tablets 10 mg

What is MAVENCLAD?

MAVENCLAD is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include relapsing-remitting disease and active secondary progressive disease, in adults. Because of its safety profile, MAVENCLAD is generally used in people who have tried another MS medicine that they could not tolerate or that has not worked well enough.

MAVENCLAD is not recommended for use in people with clinically isolated syndrome (CIS).

It is not known if MAVENCLAD is safe and effective in children under 18 years of age and is therefore not recommended.

IMPORTANT SAFETY INFORMATION

MAVENCLAD may cause serious side effects, including:

- **Risk of cancer (malignancies).** You should follow healthcare provider instructions about screening for cancer.
- **MAVENCLAD may cause birth defects if used during pregnancy.** Females must not be pregnant when they start treatment with MAVENCLAD or become pregnant during MAVENCLAD dosing and within 6 months after the last dose of each yearly treatment course. You should stop treatment with MAVENCLAD and contact your healthcare provider right away if you become pregnant during treatment with MAVENCLAD.
  - For females who are able to become pregnant:
    - Your healthcare provider should order a pregnancy test before you begin your first and second yearly treatment course of MAVENCLAD to make sure that you are not pregnant.

Please see Prescribing Information and Medication Guide, including **serious side effects**.

  - Use effective birth control (contraception) on the days on which you take MAVENCLAD and for at least 6 months after the last dose of each yearly treatment course.
  - Talk to your healthcare provider if you use oral contraceptives (the "pill").
  - You should use a second method of birth control on the days on which you take MAVENCLAD and for at least 4 weeks after your last dose of each yearly treatment course.
- For males with female partners who are able to become pregnant:
  - Use effective birth control (contraception) during the days on which you take MAVENCLAD and for at least 6 months after the last dose of each yearly treatment course.

Do not take MAVENCLAD if you:

- have cancer (malignancy).
- are pregnant, plan to become pregnant, or are a woman of childbearing age or a man able to father a child and you are not using birth control.
- are breastfeeding.
- are human immunodeficiency virus (HIV) positive.
- have active infections, including tuberculosis (TB), hepatitis B or C.
- are allergic to cladribine.

Before you take MAVENCLAD, tell your healthcare provider about all of your medical conditions, including if you:

- think you have an infection.
- have taken, take, or plan to take medicines that affect your immune system or blood cells, or other treatments for MS. Certain medicines can increase your risk of getting an infection.
- have had a recent vaccination or are scheduled to receive any vaccinations. You should not receive live or live-attenuated vaccines within the 4 to 6 weeks preceding treatment with MAVENCLAD or receive these types of vaccines during your treatment with MAVENCLAD and unless directed by your healthcare provider.
- have heart failure.
- have or have had cancer.
- have liver or kidney problems.
- are breastfeeding or plan to breastfeed. It is not known if MAVENCLAD passes into your breast milk. Do not breastfeed on the days on which you take MAVENCLAD, and for 10 days after the last dose.

How should I take MAVENCLAD?

- MAVENCLAD is given as two yearly treatment courses.
- Each yearly treatment course consists of 2 treatment weeks (also called cycles) that will be about a month apart.
- Take MAVENCLAD with water and swallow whole without chewing. MAVENCLAD can be taken with or without food.
- Swallow MAVENCLAD right away after opening the blister pack.
- Your hands must be dry when handling MAVENCLAD and washed well with water afterwards.
- Limit contact with your skin. Avoid touching your nose, eyes and other parts of the body. If you get MAVENCLAD on your skin or on any surface, wash it right away with water.
- Take MAVENCLAD at least 3 hours apart from other medicines taken by mouth during the 4- to 5-day MAVENCLAD treatment week.

- If you miss a dose, take it as soon as you remember on the same day. If the whole day passes before you remember, take your missed dose the next day. Do not take 2 doses at the same time. Instead, you will extend the number of days in that treatment week.

Your healthcare provider will continue to monitor your health during the 2 yearly treatment courses, and for at least another 2 years during which you do not need to take MAVENCLAD. It is not known if MAVENCLAD is safe and effective in people who restart MAVENCLAD treatment more than 2 years after completing 2 yearly treatment courses.

MAVENCLAD can cause serious side effects. If you have any of these symptoms listed below, call your healthcare provider right away:

- **low blood cell counts** have happened and can increase your risk of infections during treatment with MAVENCLAD. Blood tests are needed before you start treatment with MAVENCLAD, during your treatment with MAVENCLAD, and afterward, as needed.
- **serious infections such as:**
  - **TB, hepatitis B or C, and shingles (herpes zoster).** Fatal cases of TB and hepatitis have happened with cladribine during clinical studies. Tell your healthcare provider right away if you get any symptoms of the following infection related problems or if any of the symptoms get worse, including: fever, aching painful muscles, headache, feeling of being generally unwell, loss of appetite, burning, tingling, numbness or itchiness of the skin in the affected area, skin blotches, blistered rash, or severe pain.
  - **progressive multifocal leukoencephalopathy (PML).** PML is a rare brain infection that usually leads to death or severe disability. Although PML has not been seen in MS patients taking MAVENCLAD, it may happen in people with weakened immune systems. Tell your healthcare provider right away if you have any new or worsening neurologic signs or symptoms. These may include: weakness on 1 side of your body, loss of coordination in your arms and legs, decreased strength, problems with balance, changes in your vision, changes in your thinking or memory, confusion, or changes in your personality.
- **liver problems.** Blood tests should be performed to check your liver before you start taking MAVENCLAD. Symptoms of liver problems may include: nausea, vomiting, stomach pain, tiredness, loss of appetite, dark urine, or your skin or the whites of your eyes turn yellow.
- **allergic reactions (hypersensitivities).** You should stop treatment and seek immediate medical attention if any signs or symptoms of allergic reactions occur. Symptoms of an allergic reaction may include: skin rash, swelling or itching of the face, lips, tongue or throat, or trouble breathing.
- **heart failure.** MAVENCLAD may cause heart failure, which means your heart may not pump as well as it should. Call your healthcare provider or go to the closest emergency room for medical help right away if you have any signs or symptoms such as shortness of breath, a fast or irregular heart beat, or unusual swelling in your body.

The most common side effects of MAVENCLAD include: upper respiratory infection, headache, and low white blood cell counts.

These are not all the possible side effects of MAVENCLAD. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Prescribing Information and Medication Guide, including **serious side effects**.

MAVENCLAD®
(cladribine) tablets 10 mg

START THE CONVERSATION ABOUT MAVENCLAD WITH YOUR HEALTHCARE PROVIDER.

Please see Important Safety Information, MAVENCLAD Prescribing Information, and Medication Guide, including **serious side effects**.

EMD SERONO

©2020 EMD Serono, Inc. All rights reserved.

US/CLA/1018/0126a(1) 03/20